

FSRH Clinical Effectiveness Unit Statement: Use of combined hormonal contraception during the Covid-19 pandemic

18 December 2020

Why is guidance necessary?

Some contraceptive providers have voiced concern about the use of combined hormonal contraception (CHC) during the Covid-19 pandemic because of increased risk of thromboembolism.

What is the risk of thromboembolism associated with Covid-19 infection?

Evidence indicates that risk of thromboembolic events is extremely high amongst individuals hospitalised with severe Covid-19 infection.[1-7] Thromboembolic risk for individuals with less severe or asymptomatic Covid-19 infection is not established.

Why would we consider avoiding use of CHC during Covid-19 infection?

There is no published evidence to directly inform risk of thromboembolism associated with *use of CHC during Covid-19 infection*. However in general, CHC use is associated with increased risk of thromboembolism and is usually avoided in the presence of additional risk factors.[8,9]

What concerns are there about stopping CHC because of Covid-19 infection or risk of Covid-19 infection?

Combined oral contraception is the most widely used contraceptive method in the UK.[10,11] It is important that CHC users do not stop their contraception unnecessarily, with resulting risk of unplanned pregnancy. It is noted that pregnancy is associated with greater risk of thromboembolic events than CHC use.

It has been suggested (based on differences in Covid-19 outcomes between men and women and between pre- and post-menopausal women) that estrogens - both endogenous and exogenous - could potentially be protective against the immune inflammatory response associated with the most serious complications of Covid-19.[12-16] There is not, however, direct evidence to support any protective effect of estrogen-containing contraceptives.

What is the evidence for outcomes with use of other contraceptive methods during Covid-19 infection?

There is no published evidence to inform health outcomes associated with use of other effective contraceptive methods prior to and during Covid-19 infection.

What does FSRH CEU recommend?

Many questions remain to be answered on this subject. FSRH CEU suggests the following pragmatic approach:-

1. Healthy individuals

Use of CHC by eligible healthy individuals should not be restricted during the Covid-19 pandemic. CHC users should (as usual) be informed about their increased risk of thromboembolism and offered the full range of suitable alternative effective contraceptive methods.

2. Individuals with asymptomatic Covid-19 infection

Current CHC users with *asymptomatic* Covid-19 infection can continue to use CHC. Some may wish to make an immediate switch to a progestogen-only pill to try to reduce thrombotic risk, but increased risk of thrombosis persists for some time after stopping CHC.

3. Individuals with symptomatic Covid-19 infection not requiring hospitalisation

Current CHC users with *symptomatic* Covid-19 infection who do not require hospital admission should be managed according to severity of illness and degree of immobility.

- ▶ Discontinuation of CHC and initiation of a progestogen-only pill (POP) should be considered. It is noted that there could be risk of pregnancy if CHC is stopped abruptly immediately after a scheduled hormone-free interval or in the first week of use after a scheduled hormone-free interval and there has been recent unprotected intercourse. Where appropriate, emergency contraception should be discussed.
- ▶ After recovery, the individual may wish to continue POP or to consider an alternative contraceptive method that is not associated with increased risk of thromboembolism. Restarting CHC may be considered when the individual is no longer systemically unwell and has regained full mobility, bearing in mind that it is not known how long elevated thromboembolic risk associated with Covid-19 infection persists.

4. Individuals with severe Covid-19 infection requiring hospitalisation

Current CHC users requiring hospital admission because of severe Covid-19 infection should discontinue CHC.

- ▶ It is noted that there could be risk of pregnancy if CHC is stopped abruptly immediately after a scheduled hormone-free interval or in the first week of use after a scheduled hormone-free interval and there has been recent unprotected intercourse. Where appropriate, emergency contraception should be discussed.
- ▶ An alternative effective contraceptive method such as a POP should, where appropriate, be initiated prior to discharge from hospital.
- ▶ After recovery, the individual may wish to continue POP or to consider an alternative contraceptive method that is not associated with increased risk of thromboembolism. Restarting CHC may be considered when the individual is no longer systemically unwell and has regained full mobility, bearing in mind that it is not known how long elevated thromboembolic risk associated with Covid-19 infection persists.

References

1. Schulman S. Coronavirus Disease 2019, Prothrombotic Factors, and Venous Thromboembolism. *Semin Thromb Hemost* 2020 Oct;46(7):772-776.
2. Malas MB, Naazie IN, Elsayed N, *et al*. Thromboembolism risk of COVID-19 is high and associated with a higher risk of mortality: A systematic review and meta-analysis. *EClinicalMedicine* 2020 Dec;29:100639.
3. Middeldorp S, Coppens M, van Haaps TF, *et al*. Incidence of venous thromboembolism in hospitalized patients with COVID-19. *J Thromb Haemost* 2020 Aug;18(8):1995-2002.

4. Lodigiani C, Iapichino G, Carenzo L, *et al.* Venous and arterial thromboembolic complications in COVID-19 patients admitted to an academic hospital in Milan, Italy. *Thromb Res* 2020 Jul;191:9-14.
5. Wichmann D, Sperhake JP, Lütgehetmann M, *et al.* Autopsy Findings and Venous Thromboembolism in Patients With COVID-19: A Prospective Cohort Study. *Ann Intern Med* 2020 Aug;173(4):268-277.
6. Cui S, Chen S, Li X, *et al.* Prevalence of venous thromboembolism in patients with severe novel coronavirus pneumonia. *J Thromb Haemost* 2020 Jun;18(6):1421-1424.
7. Klok FA, Kruip MJHA, van der Meer NJM, *et al.* Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res* 2020 Jul;191:145-147.
8. Faculty of Sexual & Reproductive Healthcare. UK Medical Eligibility Criteria for Contraceptive Use. 2016. Available at: <https://www.fsrh.org/ukmec/> [Accessed 18/12/2020]
9. Faculty of Sexual & Reproductive Healthcare. Combined Hormonal Contraception. 2020. Available at: <https://www.fsrh.org/standards-and-guidance/documents/combined-hormonal-contraception/> [Accessed 18/12/2020]
10. Clifton S, Fuller E, Philo D. National Survey of Sexual Attitudes and Lifestyles (Natsal-3). Reference tables. 2016. Available at: <https://www.natsal.ac.uk/sites/default/files/2020-11/natsal-3-reference-tables.pdf>. [Accessed 18/12/2020]
11. Office for National Statistics. Opinions Survey Report No. 41: Contraception and Sexual Health, 2008/09. 2009. Available at: <http://webarchive.nationalarchives.gov.uk/20160105160709/http://ons.gov.uk/ons/rel/lifestyles/contraception-and-sexual-health/2008-09/index.html> [Accessed 18/12/2020]
12. Mauvais-Jarvis F, Klein SL, Levin ER. Estradiol, Progesterone, Immunomodulation, and COVID-19 Outcomes. *Endocrinology* 2020 Sep 1;161(9):bqaa127.
13. Grandi G, Facchinetti F, Bitzer J, *et al.* The gendered impact of coronavirus disease (COVID-19): do estrogens play a role? *Eur J Contracept Reprod Health Care* 2020 Jun;25(3):233-234.
14. Hamzaoglu K, Erel CT. Should estrogen be used in the co-treatment of COVID-19 patients? What is the rationale? *Maturitas* 2020 Oct;140:80.
15. Breithaupt-Faloppa AC, de Jesus Correia C, Máximo Prado C, *et al.* 17 β -Estradiol, a potential ally to alleviate SARS-CoV-2 infection. *Clinics (Sao Paulo)* 2020;75:e1980.
16. Chanana N, Palmo T, Sharma K, *et al.* Sex-derived attributes contributing to SARS-CoV-2 mortality. *Am J Physiol Endocrinol Metab* 2020 Sep 1;319(3):E562-E567.

Acknowledgements

This guidance was developed on behalf of the FSRH CEU by Dr Sarah Hardman, FSRH CEU Director and Dr. Chelsea Morroni, FSRH CEU Deputy Director.

The Clinical Effectiveness Unit (CEU) was formed to support the Clinical Effectiveness Committee of the Faculty of Sexual & Reproductive Healthcare (FSRH), the largest UK professional membership organisation working at the heart of sexual and reproductive healthcare. The FSRH CEU promotes evidence based clinical practice and it is fully funded by the FSRH through membership fees. It is based in Edinburgh and it provides a members' enquiry service, evidence-based guidance, new SRH product reviews and clinical audit/research. [Find out more here.](#)